

# A Geomorphological Atlas of the Yorkshire Wolds

Derek J. Gobbett

## Introduction

In contrast to the relatively well-researched geomorphology of the unglaciated chalklands of southern England, with the exception of Lewin (1969), little has been published on the chalk landforms of the Yorkshire Wolds. Early discussion (Cole 1879, Mortimer 1885) concentrated on the most obvious and most puzzling of features, the dry valleys and their origin.

Reid (1887), recognised that surface melt-water above a frozen substrate was instrumental in eroding the now dry valleys, yet others dismissed stream erosion entirely. Cole (1879) denied that surface water was, in any way, instrumental in valley formation. He thought that during uplift the chalk had suffered linear (tectonic) disturbances along which solution of the Chalk proceeded more rapidly than elsewhere causing surface collapse and the origin of the dry valley systems. He also thought that the present form of the dry valleys was the result of recent glacial erosion and he regarded the well-rounded chalk gravels, which reach a considerable thickness in the lower parts of the valleys, as marine beaches. In a later paper (Cole 1887) he agreed with Reid (1887) in recognising the importance of frozen ground in greatly enhancing physical weathering and erosion in the short term but regarded underground solution below the dry valleys as the most effective long term agent in the formation of these valleys.

related to  
5

Mortimer (1885) also dismissed the fluvial origin of the dry valleys. He found evidence for faults running parallel to the axes of some valleys and followed Cole in thinking that these valleys developed along "fractures in the crust" where internal solution of the chalk was more pronounced. He regarded the (dolines) on Raisthorpe Wold as incipient dry valleys. Mortimer did recognise some land slipping but failed to see that many of his lines of disturbance were due to (such processes as camber, valley bulges and) slope instability rather than tectonic movement.

elongate hollows

Foster (1985, 1987a,b,) briefly discussed the development of dry valleys and considered the fan-shaped gullies to be found at the head of many dry valleys to be post-Devensian features formed by snow melt. He thought the ice lobe in the Vale of Pickering (rose up and) in places overtopped the northern Wolds escarpment as far west as Thorpe Basset Brow and interpreted Snevver Scar [TA033788] and "channels", running parallel to the contour, as glacial drainage channels. The "channels" in question are actually formed by extensive land-slipping and lie behind (rotational slides).

fracture  
elongate masses  
by wind action or erosion  
which have slid down  
the escarpment  
(c. s.).

De Boer's studies of the Wolds dip slope suggested to him the presence of an erosion surface at about 60m and another surface at about 115m above OD. These surfaces are represented by a small number of discontinuous facets. None of the specific localities given by De Boer (De Boer *et al.* 1958, p.188) show any clearly discernable break in slope on the 1:25,000 map and as De Boer himself states, "The chief difficulties in precisely defining the higher feature are also encountered in mapping the lower terrace: first the very smooth transition from the surface of the terrace to the only

*slightly more pronounced slope of the backface, with no clear break of slope, and second, the generally subdued relief away from the valleys”*

The idea of erosion surfaces was taken up by Lewin (1969) and considerably extended using a variety of analytical techniques. He concluded that erosion surfaces do exist and proceeded to map seven surfaces in the Northern Wolds and eight in the Southern Wolds which were tentatively correlated. Map data and observation in the field have led me to think that erosion surfaces do not exist on the Yorkshire Wolds and thus do not feature in this atlas. His lower surfaces, e.g. those along the Great Wold Valley, can be shown to be terraces produced by post-glacial mass movement. Lewin did not consider how the solution of the chalk during the Tertiary would have affected the surface of the Wolds. He also failed to recognise the importance of the impact of the late Pleistocene periglacial climate on valley development and mass movement. Work in southern England (Bull 1940, Paterson 1977, Williams 1980) broadly agrees with the conclusions reached here that erosion of the chalk is greatly enhanced under cold climatic conditions when intensive physical weathering produces frost shattered chalk which is rapidly eroded during the spring melt particularly under permafrost conditions.

No detailed maps of the geomorphology of the Yorkshire Wolds have been published to my knowledge. The main object of the present work is to summarise the geomorphological features on 18 maps, each covering 10 x 7.5 km on a scale of 1:50,000. These maps are based on OS maps, DTM maps, Google Earth, aerial photographs and field observations.

### **Present landforms**

The present landforms of the Yorkshire Wolds can be divided into 5 general types.

1. A gently sloping plateau which is widely but patchily preserved on the present Wolds dip slope.
2. The escarpment.
3. A major valley, the Great Wold Valley running ENE through the northern Wolds.
4. Dry valleys which form a dense array of dendritic patterns etched into the landscape and constitute at least 80% of the Wolds surface (see Lewin 1969, fig. 13)
5. Glacial melt-water overflow channels

### Plateau areas

The Chalk outcrop of the Yorkshire Wolds is shaped rather like a west facing boomerang. Around the apex the plateau surface lies above 200m culminating at 246m at Garrowby [SE821569]. The wings of the boomerang run ENE (Northern Wolds) and SSE (Southern Wolds) and along both of these the plateau surface steadily declines to about 90m. Since the plateau is bounded by convex slopes its limits have to be somewhat arbitrarily defined.

The main watershed closely follows the top of the escarpment in the Northern Wolds except for two embayments, at Wintringham [SE880730] where the watershed is lowered to 114m, and at Wharram le Street [SE864662] where it lies at 124m. These

may be relicts of former courses of a permanent river or rivers traversing the Wolds from the west. The only other scarp valley in this part of the Wolds is Sked Dale [SE966753] just south of Sherburn which lowers the watershed from 160m to 131m. From the apex of the boomerang at Leavening Brow the watershed continues SSE but here it is much indented by several scarp slope dry valley systems, the two largest of which, at Millington and Warter, push the main watershed far to the east. Towards the concave side of the boomerang plateau height steadily decreases (see Versey 1939, fig.3) and would, before the Devensian have ended in the cliff line from Bridlington to Hessle. This concave side is now covered by the edge of Devensian glacial deposits. The strike of the Chalk generally follows the shape of the boomerang. It dips in the northern Wolds towards the SSE and in the southern Wolds to the ENE at an average gradient of 1 in 50 ( $1-2^{\circ}$ ). The surface of the dip slope along the interfluvies follows these trends but with a shallower gradient of about 1 in 100 ( $<1^{\circ}$ ).

49

On the plateau there are a number of small shallow depressions in the Chalk most of which are filled with clastic sediments and are not obvious in the landscape. They were probably originally formed as dolines. The most obvious of these is a series of linear unfilled or partially filled depressions which run *en echelon* in an east-west direction across Raisthorpe Wold [SE840625] (Mortimer 1885, Lewin 1969, Gobbett in press). The pattern of these "Raisthorpe Hollows" suggests they may be related to structural features in the Chalk.

The clastic infill of dolines was formerly used as a source of clay and sand and several of these deposits have been mapped by Mortimer (1886). More recently Bray *et al.* (1981) have recorded more than 15m of sand and brickearth filling a doline at Fimber [SE885608]. Dolines also provide natural ponds around which many of the Wolds villages were sited (Hayfield and Wagner 1995, fig.2). Larger dolines may have existed in the Tertiary landscape which were modified by surface runoff during periods of cold climate in the Quaternary. The heads of many dry valleys are bowl-shaped with small radial channels which coalesce to form a typical dry valley with a flat floor and steep sides (Fig.1). Foster (1987b, fig.12) mapped two other examples near Octon Lodge [T010700]. Such bowl-shaped depressions might have originally developed as dolines on the pre-Pleistocene surface. Figure 1 shows the widespread distribution of these depressions.

Solution pipes are probably common and widespread on the plateau but are rarely visible except in chalk pits and quarries. They are usually filled with a flint-dominated breccia which may be cemented into "fairystone" seen *in situ* in Fairydale and at St Austin's Rock (Gobbett 2006). Boulders of "fairystone" are commonly found along field margins and road verges where they have been moved to avoid the damage they would do to the plough.

The plateau with its dolines and pipes and an average gradient of less than  $1^{\circ}$  is likely to have extended over the whole area of the Wolds during the long progress of Chalk erosion by internal solution during Tertiary time. Differential subsidence of the eroding Chalk could have produced broad shallow facets on the dip slope simulating poorly preserved erosion surfaces.

### The escarpment

Along the front of the Wolds boomerang the Chalk is underlain for the most part by clays. The relatively slower erosion of the chalk results in a marked escarpment the character of which varies along its length. It is convenient to divide it into three sections anticlockwise from northeast to south.

### *1. Flotmanby to Knapton*

Here the escarpment runs WSW almost in a straight line. The top rises from 120m at Flotmanby Brow to 160m at Knapton: the base lies constantly at about 50m. The base of the Chalk also rises from 50m at Flotmanby to 80m at Knapton, so although it is underlain by the Speeton Clay, the steep slope of the escarpment is formed entirely or largely of Chalk (Fig.2a). The Sherburn Sands (probably of Devensian age) overlap the base of the Chalk up to about the 70m contour. There are only two small gaps (mentioned above in describing the watershed) one at Ganton and one at Sherburn where narrow scarp (obsequent) dry valleys have cut through the escarpment top.

The inherent instability of Chalk overlying clay has caused landslips all along the escarpment. These typically break away from near the top of the escarpment along a curved fracture and move down-slope to form a buttress with its long axis parallel to the slope. In between these slips the original scarp appears as a small valley or gully. Where the Chalk has been exposed e.g. in the pit on Willerby Brow [TA012783], it shows cambering and gulls. Westwards the slipped area increases in width and in size of individual slips as the base of the Chalk rises and thus the thickness of clay in the scarp face increases. It is particularly well developed below East Heslerton Brow where there is evidence of rotational slides. One of these is well exposed in a chalk pit at SE938761 where the Lower Chalk dips 25° south.

### *2. Knapton to Leavening Brow*

From Knapton the scarp turns southwest. It is here more discontinuous with detached or partly detached outliers. At Wharram le Street and near Duggleby the escarpment consists entirely of Kimmeridge Clay.

The base of the escarpment rises from 50m to 120m but as the base of the Chalk also rises from 80m to 180m it forms an increasingly thin capping (Fig.2b). As a result there are extensive landslips which are particularly well developed on Birdsall Brow. The Stack Hills [SE8771] on the south side of the Wintringham Gap are deep rotational slips in which the Chalk dips 70° south towards the escarpment. In the larger Chalk outliers of Wintringham, Thorpe Basset, Duggleby, Settrington and Grimston Brow the Chalk appears to be tilted probably by cambering and some anomalously high dips are marked on the BGS map. Above North Grimston is a secondary escarpment formed of Corallian limestone and separated from the Chalk by a thin layer of Kimmeridge Clay.

### *3. Leavening Brow to North Ferriby*

This 40km stretch runs SSE, overall in a straight line, but it is broken by over 30 dry valleys which form the heads of spring line streams. These range from single coombes e.g. Cleaving Coombe [SE860465] to extensive basins like those around Millington and Warter which rival in size some of the dip slope drainage basins.

The top of this part of the escarpment falls from 200m to 90m and its base from 120m to 20m. The base of the Chalk drops from 180m to sea level (Fig.2c). The strata

underlying the chalk are more variable but Lias clays predominate overlain in places by thin Middle Jurassic and Oxfordian strata.

The Chalk escarpment remains unstable and features landslips but generally on a smaller scale than those further north. At Partridge Hall Quarry [SE858470] the Chalk dips eastward into the escarpment at 15-20° and is cut by numerous normal faults striking parallel to the trend of the escarpment and dipping 70-80° west. These faults slice up the chalk to produce a series of rotational blocks caused by gravity collapse over the underlying clays. The Black band is exposed at the base of the quarry and this also may have provided a weak layer over which the Chalk slipped (R. Myerscough, personal communication). copy

Flanking the entrance to the Millington basin thin limestone bands in the Lower Lias form a low wooded secondary escarpment below the Chalk.

### The Great Wold Valley

From its head at Wharram le Street the Great Wold Valley runs ENE to Burton Fleming beyond which it abruptly turns south to Rudston. Here the present valley turns sharply through another right angle and runs to the sea at Bridlington Quay. This part of the valley narrows and its longitudinal gradient steepens. The Great Wold Valley drainage basin is many times larger than any other basin on the Wolds.

In comparison with the largest dip slope dry valley the Great Wold Valley is broader, has a less steep longitudinal gradient and more gentle sides which, although commonly modified by landslips, tend to be concavo-convex in contrast to the convex slopes of dry valleys (Fig.3).

Along the base of the valley sides a number of small lobe-like rounded hills intrude into the valley floor. These occur mainly on the south, north-facing side of the valley and increase in size down valley. These small hills are usually cultivated and show no exposures of the Chalk. Their contours are smoother than those of the rotational slides of the Stack Hills on the south side of the Wintringham Gap. They appear to have been formed by mass movement of a regolith of brecciated chalk which has denuded the upper part of the slope and aggraded the lower part. This type of mass movement is probably responsible for a terrace seen in some places on the lower part of the valley side.

Although its tributary valleys are dry the main valley carries some surface water. At the present time springs from the base of the Chalk feed a small stream which has a permanent flow from Wharram le Street to Kirby Grindalythe. Between Kirby Grindalythe and West Lutton stream flow relies on a rise in the water table and is thus intermittent. Below West Lutton only after periods of exceptionally high rainfall does water rise through the valley floor to produce a surface flow which may continue as far as Butterwick before sinking into the ground. More commonly ground water rises between Foxholes and Wold Newton from where a broader stream floods the valley floor. At Rudston a permanent spring sources the stream to the coast at Bridlington Quay.

This typical "winterbourne" known locally as the "Gypsey Race" is clearly a misfit. "Gypsey" is pronounced with a hard "G". Its origin is obscure but it may come from the Norse "gypa" which means a gushing spring. Taylor (1996) notes that it's earliest



mention is found in the chronicle of William of Newburgh who was born in Bridlington in 1136. He wrote "*Famosae illae aquae quos volgo Gipse vocant*" (This famous watercourse called Gipse in the vulgar tongue). Allen (1831) provides an excellent description.

*"The Gipseys are streams of water which at different periods are observed on some parts of the Yorkshire wolds; they appear towards the latter end of winter or early in the spring. They are then seen tricking through the grass where the ground is not broken, and sometimes rushing with considerable force through the surface; and the emission of water is often so copious as to constitute a very considerable stream, filling a drain twelve feet wide and three feet deep, called by the country-people the gipsey-race, ... .. it is undoubtedly accelerated and augmented by a continuance of heavy rains; and indeed the gipseys never make their appearance except in a very wet season, when they sometimes flow during two or three months, and then totally cease, leaving scarcely a mark to distinguish the place from which the water issued"*

#### 1.4 The Dry Valleys

The Wolds plateau is dissected by numerous valleys which have all the characteristics of a fluvial origin but are presently dry. The valley sides are smooth, rounded, convex slopes which are mantled, as are the remains of the original plateau and the valley floors, by a regolith of brecciated chalk with a matrix of loess (Catt *et al.* 1974, Neal 2011).

The valleys display a dendritic pattern in which the tributaries often join the main valley at high angles approaching  $90^{\circ}$ . This led Lewin (1969) to consider that they might be joint controlled. Evidence of joint patterns in the Wolds Chalk is limited and what is known suggests they are complex. High confluence angles can be explained by the fact that where they occur valley sides are steep,  $>20^{\circ}$ . On the dip slope the dry valleys are organised into a number of small drainage basins which are elongated parallel to the plateau slopes and have narrow exits. On the scarp slope Lewin (1969 fig. 13) maps 38 basins but 28 of these are small simple valleys with very small catchments.

Typically the head of a dry valley is a gentle depression traversed by several shallow gullies which converge. Down valley the gradient steepens markedly and becomes entrenched into the plateau. Unless they have been modified by landslips, the valley sides usually slope at  $22^{\circ}$ . This angle is remarkably constant all over the Wolds and is only occasionally exceeded up to an angle of  $30^{\circ}$ . The slopes are convex; the flat valley floor has a steep longitudinal gradient and in places shows traces of small channels separated by low ridges. Small tributary valleys running at right angles to the main valley commonly show signs of a small alluvial fan at their mouths.

The majority of the dry valleys have no marked asymmetry in cross section but some 30 valleys do clearly show an asymmetry. In these valleys most of the steeper slopes face between WNW and NNW as shown in Figure 4. Cole (1879) also noted that the steeper sides of the Wold valleys "as a rule" face north and west. Ollier and Thomasson (1957) who studied asymmetrical dry valleys in the Chilterns found "that in most cases where a valley is asymmetrical it is the west facing slopes and to a lesser extent the south facing slopes that are the steeper". It is difficult to explain the dominance of the NW-facing steeper slopes, as in a cold climate the west and south facing slopes may be regarded as thawing more and thus suffering more rapid erosion.

In plan the valleys are quite variable. Regular interlocking spurs are common but many valleys are quite straight or have straight sections. The sides may be symmetrical or asymmetrical and the orientation of the asymmetry is not constant.

In plan the valleys are quite variable. Regular interlocking spurs are common but many valleys are quite straight or have straight sections. The sides may be symmetrical or asymmetrical and the orientation of the asymmetry is not constant. Down valley the longitudinal gradient decreases, the sides become less steep and the flat floor widens progressively to 200m or more. It is now underlain by well rounded chalk gravel with angular flints best exposed by Station Cottages just north of Wetwang [SE935595] and in Garton Slack [SE959596] where it has been extensively quarried in the past.

The floor

The steep sides of dry valleys commonly show signs of landslips. These are similar to but smaller in scale than the slips of the escarpment. Another common feature is a marked terrace along the base of the valley side raised a couple of metres above the valley floor, e.g. at Hog walk [SE916658] and Lang Dale [TA046774]

A narrow gully running straight down the steep side of a dry valley is not uncommon. This erosive feature which may show the remains of a small debris fan at its base I have termed a *Torrent Scar*. Torrent scars have been observed to be eroded almost instantaneously during periods of very heavy rainfall when ground water bursts out of the hillside and cuts a narrow gully sometimes two to three metres deep through the brecciated chalk regolith leaving a deposit of angular chalk debris over the valley floor (Hood 1892, Marley 1991). Subsequent collapse of the gully sides and re-growth of vegetation leave a scar down the valley side. Snevver Scar which Foster (1987c) regarded as a glacial meltwater channel, may be an example of a large torrent scar.

### Glacial melt water overflow channels

On the lower part of the dip slope adjacent to the edge of the glacial drift some interfluves have been cut across by sinuous channels with flat floors and concave sides, bowl-shaped in cross section. These channels were described by De Boer (1945) who interpreted them as overflow channels draining a series of proglacial lakes which flooded the ends of the Wolds dip slope valleys during the summers of the late Devensian Dimlington Stadal. Figure 5 is an interpretation of the Wolds dip slope at that time.

The channel which occupies Hunmanby Dale [TA100750] slopes southwards from an altitude of 70m to 30m and could have carried melt-water from the ice lobe at the east end of the Vale of Pickering into a proglacial lake flooding the lower reaches of the Great Wold Valley. All the valleys draining eastwards from the Wolds would have been blocked by the ice front which would not have been static and would also have supplied copious amounts of melt water.

Although some water may have leaked out under the ice in sub-glacial channels e.g. the channels at Risby Park [TA000350], a series of proglacial lakes which at any one time probably differed in the altitude of their shorelines would have occupied the lower part of these valleys. By far the largest drainage area is that of the Great Wold Valley and the melt-water from this combined with that from the Pickering ice lobe produced the largest lake, filling the lower reaches of the Great Wold Valley, which we may call Lake Rudston. If the altitude of the lake surface increased to 50m water could have spilled over the Kilham Lane col (48m) into another lake at Kilham. The next col to the southwest lies at 64m so the lake waters must have risen further to be able to spill over into the next lake at Garton. South of Garton the low ground is now largely drift covered and any overflow channels may lie underneath the drift deposits.

Between North Dalton and Middleton the interfluvium is cut across by a channel rising to 63 m. South of Middleton a sinuous channel drained south to Goodmanham Dale where it cut across the watershed, lowered it to 50m and flowed down Spring Dale and into the Vale of York and Lake Humber.

### Evolution of the landforms

Unlike the chalklands of southern England no Tertiary sediments have been reliably recorded from the Yorkshire Wolds and the history of uplift since the end of the Cretaceous is obscure. Monoclinial folding and faulting of the Chalk, produced by fault movements in the underlying Jurassic basement are probably of early to mid Tertiary in age (Starmer 2008). These structures trend east-west but show little direct relationship to any of the main geomorphological features.

The Great Wold Valley is a major element of the landscape and was probably formed by a relatively large pre-Quaternary river rising far to the west and traversing the Wolds possibly through the Winteringham Gap. It may have been comparable with the rivers that currently cut through the North and South Downs of southern England.

The processes of weathering and denudation of the Chalk are climate dependent. In the temperate climate which prevailed during much of the Cenozoic slow chemical weathering would have predominated and internal solution of the Chalk would have lowered the surface leaving a regolith of clay-with-flints. Internal solution of the Chalk would have taken place most probably beneath a forest cover and would not have been uniform. The subsiding surface would have been diversified by enclosed hollows (dolines) and possibly by steps across the main dip slope simulating superficial "erosion surfaces". To get some idea of erosion rates, if it took 60 million ( $6 \times 10^7$ ) years to remove 600m ( $6 \times 10^8$   $\mu\text{m}$ ) of chalk the average rate of solution would be  $10 \mu\text{m yr}^{-1}$ . This may be compared to Cole's 1879 estimate of one foot in 5779 years (c.  $50 \mu\text{m yr}^{-1}$ ). Kerney *et al.* (1963, p.188) refer to Atkinson (1957) "*who presents excellent evidence from archaeological excavations on the Chalk to show that a lowering of between 15 and 20 in. on level ground during the last 4000 years is by no means uncommon*". This equates to 9.5 to  $12.5 \mu\text{m yr}^{-1}$ .

The marked changes in climate during the Pleistocene introduced several cold periglacial periods with intervals of permafrost. Under these conditions the erosion of the Chalk would have been greatly accelerated. Physical weathering by summer freeze and thaw and erosion by mass movement and melt water could have eroded the Chalk surface perhaps an average of 1mm to 1cm per year which is  $10^3$  to  $10^4 \mu\text{m yr}^{-1}$ . Thus recent periglacial denudation could have proceeded from one hundred to one thousand times faster than the temperate denudation by chemical solution and thus left a far more obvious footprint on the present landscape. Kerney *et al.* (1963) show that coombes on the escarpment of the North Downs in Kent were almost entirely eroded by solifluction and melt water in about 500 years during the Late Glacial, in the cold phase of pollen zone III when the ground remained frozen in winter.

The Yorkshire Wolds have been overridden by an ice sheet at least once during the Pleistocene and it is reasonable to assume that the ice removed the clay-with-flints regolith although some small patches remain (Matthews 1977). The present aspect of the landforms is largely due to the weathering and erosion that occurred during the late Devensian when the whole area must have been subjected to permafrost. The now



dry valleys appear as fresh young features little modified during the Holocene. The slope angle of the valley sides (around 22°) is consistent with the angle of rest of chalk scree. The steep longitudinal gradient and flat floor typical of the valleys suggests that they were occupied by intermittent flashy braided streams carrying a gravel bed load in the Devensian summer.

During the early post-glacial gradual thaw of the permafrost would have encouraged mass movement to produce the extensive landslips and rotational slides which now diversify the Chalk escarpment, the sides of the Great Wold Valley, and to a lesser extent the dry valleys.

## References

- Allen, T. 1831. *New and complete history of the county of York*, vol.4.
- Atkinson, R.J.C. 1957. Worms and weathering. *Antiquity*, **33**, 219-233
- Bray, D.A., 1981. Quaternary sediments at Westfield Farm, Fimber. *Proc. Yorks. geol. Soc.*, **43**, 377-393
- Bull, A.J. 1940. Cold conditions and landforms in the South Downs. *Proc. geol. Assoc.* **51**, 63-71.
- Catt, J. A., Weir, A.H. and Madgett, P.A. 1074. The loess of eastern Yorkshire and Lincolnshire. *Proc. Yorks. geol. Soc.* **40**, 23-29
- Cole, E.M. 1879. On the origin and formation of the Wold dales. *Proc. Yorks. geol. Soc.* **7**, 128-140
- De Boer, G. 1945. A system of glacier lakes in the Yorkshire Wolds. *Proc. Yorks. geol. Soc.* **25**, 223-33
- De Boer, G., Neal, J.W. and Penny, L.F. 1958. A guide to the geology of the area between Market Weighton and the Humber. *Proc. Yorks. geol. Soc.*, **31**, 157-209
- Ellis, S. (ed) 1987 *East Yorkshire field Guide*. Quat. Res. Assoc. London.
- Foster, S.W. 1985. *The late glacial and early postglacial history of the Vale of Pickering and northern Yorkshire Wolds*. PhD thesis: Univ. of Hull
- Foster, S.W. 1987a. The Sherburn Sands of the southern Vale of Pickering. In Ellis, S. 1987, pp. 31-35.
- Foster, S.W. 1987b. The dry drainage system on the Yorkshire Wolds. In Ellis, S. 1987, pp.36-38
- Foster, S.W. 1987c. Features of glacial drainage on the northern Wolds escarpment. In Ellis 1987, pp.39-41.
- Gobbett, D.J. 2006. St Austin and the Fairy: a tale of two RIGS. *Humberside Geologist* **14**, ~~48-68~~. 33-38
- Gobbett, D. J. (In press) A summary Cenozoic history of the Yorkshire Wolds, *Humberside Geologist*, **15**, ~~48-58~~.
- Hayfield, C. and Wagner, P. 1995. From dolines to dew ponds: a study in water supplies on the Yorkshire Wolds. *Landscape History*, **17**, 49-65
- Hood, J.D. 1892. *Waterspouts on the Yorkshire Wolds: cataclysm at Langtoft and Driffield*. Driffield: Frank Fawcett
- Kerney, M.P., Brown E.H. and Chandler, T.J. 1963. The late glacial and postglacial history of the Chalk escarpment near Brook, Kent. *Phil. Trans. Roy. Soc. (B)*, **248**, 135-204
- Lewin, J. 1969. The Yorkshire Wolds, a study in geomorphology. *Univ. Hull. Occas. Pap. Geogr.*, **11**, 89p

- Marley, D.S. 1991. *Souvenir of the great flood at Driffield, Friday 20<sup>th</sup> May 1910*. Reprinted from the "Driffield Observer"
- Matthews, B.G. 1977. Clay-with-flints on the Yorkshire and Lincolnshire Wolds. *Proc. Yorks. geol. Soc.* **41**, 231-9
- Mortimer, J.R. 1885. On the origin of the Chalk Dales of Yorkshire. *Proc. Yorks. geol. Soc.* **9**, 29-42.
- Mortimer, J.R. 1886. *The surface geology and modern enclosure of Fimber on the Yorkshire Wolds*. Unpublished maps. Museum of transport and Hydrology, Hull.
- Neal, C. 2011. *People and the environment: a geoarchaeological approach to the Yorkshire Wolds landscape*. PhD thesis: Univ. of York
- Ollier, C.D. and Thomasson, A.J. 1957, Asymmetrical valleys of the Chiltern Hills. *Geogr. J.* **123**, 71-80
- Paterson, K. 1977. Scarp face dry valleys near Wantage, Oxfordshire. *Trans Inst. Br. Geog. n.s.* **2**, 192-204
- Reid, C, 1887. On the origin of dry chalk valleys and of coombe rock. *Quart. J. geol. Soc. Lond.* **43**, 364-373
- Starmer, I.C. 2008. The concentration of folding and faulting in the Chalk at Staple Newk (Scale Nab) near Flamborough, East Yorkshire. *Proc. Yorks. geol. Soc.* **57**. 95-106
- Taylor, J.F. 1996. *A history of Butterwick and Boythorpe*. Foxholes
- Versey, H.C. 1939. The Tertiary history of East Yorkshire. *Proc. Yorks. geol. Soc.* **23**, 302-214
- Williams, R.B.G. 1981. The weathering and erosion of chalk under periglacial conditions. In Jones, D.C.K. *The geomorphology of the British Isles. Southeast and southern England*, pp.225-248

A Geomorphological Atlas of the Yorkshire Wolds  
by Derek Gobbett

Copyright - Hull Geological Society 2020